



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## SOLUTIONS OF EXERCISES.

## 174

FIND the locus of the intersection of the altitudes of a triangle whose base and area are given.

SOLUTION.

Let the extremities of the base be  $(\pm c, 0)$  and the vertex  $(x, h)$ .  
Then the declivity of the side is

$$m = \frac{h}{x - c},$$

and the declivity of the perpendicular to it

$$m' = \frac{y}{x + c}.$$

The condition of perpendicularity gives

$$hy = c^2 - x^2,$$

which represents a parabola with vertical axis passing through the extremities of the base.  
[*H. B. Newson; T. U. Taylor.*]

## 180

FIND the envelope of a system of circles each of which is seen from two fixed points under a constant angle.

SOLUTION.

Let the line joining the fixed points be the  $x$ -axis and its middle point be the origin. Let  $2c$  be the distance between the points; let  $\sin^{-1}(m)$  be half the constant angle.

It is evident that the locus of the centres of the system is the  $y$ -axis. Let  $(0, a)$  be the co-ordinates of the centre of any circle of the system. Then

$$r^2 = (a^2 + c^2)m^2,$$

and the equation to the circle is

$$x^2 + (y - a)^2 = (a^2 + c^2)m^2.$$

Differentiating with respect to the parameter  $a$ , we obtain

$$-2(y - a) = 2am^2;$$

whence

$$a = \frac{y}{1 - m^2}.$$

This gives for the equation to the envelope

$$\frac{x^2}{m^2} - \frac{y^2}{1 - m^2} = c^2,$$

which represents an hyperbola with the fixed points as foci and  $e = m^{-1}$ .

[*H. B. Newson.*]

### 228

SUPPOSE a vessel in the form of an inverted frustum of a cone to be filled with water, and then tipped until as much water runs out as is possible without uncovering any part of the bottom; required, the volume of the water which remains.

SOLUTION.

Let  $R, r$  be the radii of the bases of the frustum,  $H$  altitude,  $a, b$  semi axis of the elliptic water surface. The supplemental cone has a volume

$$v = \frac{1}{3} \pi H \frac{r^3}{R - r}.$$

This added to the required volume gives an oblique cone whose altitude is

$$k = \frac{2hRr}{a(R - r)},$$

and whose volume, therefore, is

$$\frac{1}{3} \pi abk = \frac{1}{3} \pi H \cdot \frac{b^3}{R - r}.$$

Hence the required volume is

$$V = \frac{1}{3} \pi H \cdot \frac{b^3 - r^3}{R - r},$$

in which

$$b = \sqrt{Rr},$$

as may be easily proved.

[*B. E. Sheppard.*]

Also solved by *Mr. O. L. Mathiot.*

### 230

SOLVE the equations

$$x^2(y - z) = a,$$

$$y^2(z - x) = b,$$

$$z^2(x - y) = c.$$

[*Frank Morley.*]

SOLUTION.

$$a + b + c = -(y - z)(z - x)(x - y),$$

$$abc = x^2 y^2 z^2 (y - z)(z - x)(x - y),$$

whence  
is known.

$$xyz = k$$

Accordingly

$$ayz = k(xy - zx),$$

$$bzx = k(yz - xy);$$

whence

$$ayz + bzx = k(yz - zx),$$

or

$$\frac{x}{k - a} = \frac{y}{k + b},$$

$$\frac{y}{k - b} = \frac{z}{k + c},$$

$$\frac{z}{k - c} = \frac{x}{k + a},$$

are obtained in the same way, By means of these relations we eliminate  $y$  and  $z$  (say) from the first of the given equations and find  $x$ ; thus

$$x^3 \left[ \frac{k + b}{k - a} - \frac{k - c}{k + a} \right] = a.$$

The value of  $k$  is

$$k = \sqrt{\frac{-abc}{a + b + c}}$$

[*T. U. Taylor.*]

## 231

IF we take products of  $n$  consecutive terms of the arithmetical series  $a, a - d$ , etc., commencing for the first product with the first term, for the second product with the second term, and so on; and then multiply these products by the coefficients in the expansion of  $(1 - x)^n$ , the aggregate will be  $n!d^n$ , which is independent of the first term. [*W. W. Johnson.*]

SOLUTION.

If the  $n$  differences of a series of  $n + 1$  quantities be multiplied by the coefficients in the expansion of  $(1 - x)^{n-1}$ , the aggregate is readily seen to be the same as that of the  $n + 1$  quantities multiplied by the coefficients of  $(1 - x)^n$ .

Now the first product named in the question is

$$a(a - d) \dots [a - (n - 1)d],$$

and the second is

$$(a - d) \dots [a - (n - 1)d] (a - nd).$$

Thus their difference is

$$nd \times (a - d) \dots [a - (n - 1)d].$$

Considering in like manner the differences between each pair of consecutive products, the theorem above shows that the aggregate in question is the product of  $nd$  into a similar aggregate in which  $n$  is changed to  $n-1$ , and  $d$  is unchanged. Thus denoting the aggregate by  $S_n$ , we have

$$S_n = nd \cdot S_{n-1} = nd(n-1)d \cdot S_{n-2} = \dots = n!d^n,$$

since  $S_1 = d$ .

[W. W. Johnson.]

Also solved by *Dr. R. A. Harris*.

### 234

IN a regular heptagon  $ABCDEFG$ , show that

$$\frac{1}{AB} = \frac{1}{AC} + \frac{1}{AD}. \quad [\text{Frank Morley.}]$$

SOLUTION.

Let  $x = \frac{1}{7}\pi$ ; then

$$\begin{aligned} \frac{1}{\sin x} - \frac{1}{\sin 3x} &= \frac{2 \cos 2x \sin x}{\sin x \sin 3x} \\ &= \frac{2 \cos 2x}{\sin 4x} \\ &= \frac{1}{\sin 2x}, \end{aligned}$$

which was to be proved.

[T. U. Taylor.]

Also solved by *Dr. R. A. Harris*.

### 235

FIND the areas of the greatest and of the least rhombus inscribed in an ellipse. [R. H. Graves.]

SOLUTION I.

The central radius vector of an ellipse is given by the formula

$$\begin{aligned} \frac{1}{r^2} &= \frac{\cos^2 \theta}{a^2} + \frac{\sin^2 \theta}{b^2} \\ &= \frac{1}{b^2} (1 - e^2 \cos^2 \theta). \end{aligned}$$

If  $r_1, r_2$  be orthogonal radii, the area of the rhombus of which they are semi-diagonals is

$$2r_1r_2 = \frac{2b^2}{\sqrt{[(1 - e^2 \cos^2 \theta)(1 - e^2 \sin^2 \theta)]}}.$$

This has its least value when

$$(1 - e^2 \cos^2 \theta)(1 - e^2 \sin^2 \theta) = z$$

is greatest, and its greatest value when this is least. But

$$z = 1 - e^2 + \frac{1}{4}e^4 \sin^2 2\theta$$

is greatest when  $\sin 2\theta = 1$ , and least when  $\sin 2\theta = 0$ . The corresponding values of the area are

$$Q' = \frac{2b^2}{1 - \frac{1}{2}e^2}, \quad Q'' = \frac{2b^2}{\sqrt{1 - e^2}};$$

or 
$$Q' = \frac{4a^2b^2}{a^2 + b^2}, \quad Q'' = 2ab. \quad [W. M. Thornton.]$$

#### SOLUTION II.

Let the ellipse be projected into a circle whose diameter is equal to the minor axis. Every parallelogram inscribed in the ellipse becomes a rectangle inscribed in the circle. In case of the rhombus, the sides being as 1 : 1, after projection the ratio of two adjacent sides of the rectangle may vary from 1 to  $\frac{b}{a}$ . Find now the greatest and least rectangles the ratios of whose adjacent sides lie within these limits.

The first is a square. For, draw axes through the centre of the rectangle parallel to its sides (the  $x$ -axis being parallel to the longer side), and let  $\theta$  denote the angle which the diagonal makes with the  $x$ -axis. Then the area is

$$u = b^2 \sin \theta \cos \theta.$$

$$\therefore \frac{du}{d\theta} = b^2 \cos 2\theta;$$

and  $\theta = 45^\circ$  causes  $\frac{du}{d\theta}$  to vanish.

We can project this square into a rhombus only by projecting the diagonals into the major and minor axes. Hence, the area of the greatest rhombus is  $2ab$ .

$u$  being an increasing function up to  $\theta = 45^\circ$ , the smaller we can take  $\theta$ , the smaller will be the area. This value is  $\theta = \tan^{-1} \left( \frac{b}{a} \right)$ ; i. e. the ratio of the two adjacent sides of our rectangle is  $\frac{b}{a}$ . Hence, the length of its longer side is  $\frac{2ab}{\sqrt{a^2 + b^2}}$ .

We can project this rectangle into a rhombus only by projecting the shorter sides into lines parallel to the major axis; the longer sides into lines parallel to the minor. This rhombus, then, is a square whose area is  $\frac{4a^2b^2}{a^2 + b^2}$ .

[R. A. Harris.]